

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/261376462>

Effect of eplerenone in percutaneous coronary intervention-treated post-myocardial infarction patients with left...

Article in *European Journal of Heart Failure* · June 2014

DOI: 10.1002/ejhf.88

CITATIONS

3

READS

49

8 authors, including:



Renaud Fay

Centre Hospitalier Universitaire de Nancy

66 PUBLICATIONS 1,342 CITATIONS

[SEE PROFILE](#)



Julian Gunn

The University of Sheffield

55 PUBLICATIONS 984 CITATIONS

[SEE PROFILE](#)



Bertram Pitt

University of Michigan

563 PUBLICATIONS 37,885 CITATIONS

[SEE PROFILE](#)



Faiez Zannad

Centre Hospitalier Universitaire de Nancy

876 PUBLICATIONS 53,206 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Imaging and physiology assesment of CAD [View project](#)



Acute Coronary Syndromes [View project](#)

Effect of eplerenone in percutaneous coronary intervention-treated post-myocardial infarction patients with left ventricular systolic dysfunction: a subanalysis of the EPHESUS trial

Javaid Iqbal^{1*}, Renaud Fay², David Adlam³, Iain Squire³, Yasir Parviz¹, Julian Gunn¹, Bertram Pitt⁴, and Faiez Zannad²

¹Department of Cardiovascular Science at the University of Sheffield, and Cardiology Department at Sheffield Teaching Hospitals NHS Trust, Sheffield, UK; ²INSERM, Centre d'Investigation Clinique and Centre Hospitalier Universitaire, and the Department of Cardiology, Nancy University, Université de Lorraine, Nancy, France; ³Department of Cardiology, and NIHR Cardiovascular Biomedical Research Unit, Glenfield Hospital, Leicester, UK; and ⁴Cardiovascular Centre, University of Michigan, Ann Arbor, MI, USA

Received 27 December 2013; revised 14 February 2014; accepted 21 February 2014; online publish-ahead-of-print 4 April 2014

Aims	EPHESUS was a multicentre, double-blind clinical trial in which 6632 patients with acute myocardial infarction (AMI) complicated by LV systolic dysfunction (LVSD) were randomized to receive eplerenone ($n = 3319$) or placebo ($n = 3313$). A total of 1580 EPHESUS patients were treated with PCI, which is now the standard treatment for AMI. This EPHESUS substudy examined the effects of eplerenone upon cardiovascular outcomes in PCI-treated patients.
Methods and results	EPHESUS patients were divided into PCI-treated and non-PCI-treated cohorts, and the effect of eplerenone upon mortality and other major adverse cardiovascular outcomes was assessed in each cohort. The PCI-treated patients ($n = 1580$) were younger, and had better renal function and fewer co-morbidities than non-PCI-treated patients ($n = 5052$). Cardiovascular mortality was significantly lower in PCI-treated patients as compared with non-PCI-treated patients (7% vs. 16%, $P < 0.0001$). However, the incidence of non-fatal events was similar in PCI-treated and non-PCI-treated cohorts. There was no statistical difference between the PCI-treated and non-PCI-treated cohorts in the primary or secondary outcomes of the trial. Eplerenone administration, compared with placebo, in the PCI-treated cohort did not affect PCI-related clinical outcomes, including recurrence of angina, the occurrence of acute coronary syndromes, or the need for further revascularization.
Conclusions	The beneficial effects of eplerenone in the EPHESUS trial exist for both PCI- and non-PCI-treated AMI patients with LVSD. Eplerenone has minimal, if any, effect upon reducing PCI-related adverse events in the PCI-treated cohort.
Keywords	Eplerenone • Heart failure • Angioplasty • Myocardial infarction

Introduction

Acute myocardial infarction (AMI) and consequent LV systolic dysfunction (LVSD) are the leading causes of morbidity and mortality. Mineralocorticoid receptor (MR) antagonism with eplerenone has

been shown to improve survival and reduce hospitalization due to cardiovascular events in patients with AMI and LVSD in the landmark EPHESUS (Eplerenone Post-Acute Myocardial Infarction Heart Failure Efficacy and Survival Study) trial.¹ In this trial, a mean dose of 43 mg of eplerenone produced a significant reduction in

*Corresponding author: M116, Department of Cardiovascular Science, University of Sheffield, Beech Hill Road, Sheffield S10 2RX, UK. Tel: +44 114 2268986, Fax: +44 114 2711863, Email: javaid@doctors.net.uk

all-cause and cardiovascular mortality compared with placebo.¹ National and international guidelines therefore strongly recommend using MR antagonists for patients with post-infarct LVSD.^{2,3}

In contemporary practice, patients with AMI are usually treated with PCI, involving balloon angioplasty and stent implantation.^{4,5} However, only a quarter of patients in the EPHESUS trial received PCI for treatment of AMI while others were treated with thrombolysis, coronary artery bypass grafting (CABG), or conservative medical treatment.¹ Therefore, application of the EPHESUS data to current practice can be questioned. We aimed to evaluate the impact of eplerenone administration in heart failure among patients managed with PCI.

Experimental data have suggested that eplerenone may improve the PCI outcomes by reducing neointimal proliferation⁶ and accelerating endothelial regeneration.^{7,8} While pre-clinical studies support these hypotheses,^{9–11} there has been no study investigating this potential benefit of eplerenone in humans. Although the EPHESUS trial did not have a pre-specified aim to examine PCI outcomes, and there was no routine angiographic follow-up, surrogate clinical markers were systematically recorded. The second objective of this study was to evaluate the impact of eplerenone upon PCI-related adverse clinical outcomes, including recurrence of angina, the occurrence of acute coronary syndromes, and the need for repeat revascularization.

Methods

EPHESUS trial

EPHESUS was a large multicentre, double-blind randomized controlled clinical trial that assessed the impact of eplerenone upon clinical outcomes in patients with LVSD after AMI.¹ Patients ($n = 6632$) were randomized to receive eplerenone ($n = 3319$) or placebo ($n = 3313$). Eplerenone was initiated at 25 mg/day at ~7 days after AMI and titrated at 4 weeks to 50 mg/day, if serum potassium was <5 mmol/L. About 24% patients in the EPHESUS trial received PCI as a treatment of AMI while others were treated with thrombolysis (27%), CABG (1%), or conservative medical treatment (48%). Patients were followed-up for an average of 16 months.

EPHESUS PCI substudy

This substudy evaluated the effect of eplerenone administration in the PCI-treated subgroup of patients in the EPHESUS trial. We analysed the PCI-treated ($n = 1580$) vs. non-PCI-treated ($n = 5052$) patients. Furthermore, we compared the baseline characteristics and clinical outcomes in PCI-treated patients in the eplerenone- ($n = 799$) and placebo- ($n = 781$) treated groups.

Statistical analysis

Data are presented as mean $\pm$ SD or proportion and percentage, as indicated. Continuous variables were analysed using the Mann–Whitney test and categorical variables with the χ^2 test. Multivariate analyses were performed using the Cox regression method. Hazard ratios (HRs) along with the 95% confidence interval (CI) are presented. P -values <0.05 were considered significant. All statistical analyses were performed using SAS[®] 9.2 software (SAS Institute, Cary, NC, USA).

Results

The EPHESUS PCI cohort is different from the non-PCI cohort

The baseline demographic and clinical characteristics of PCI-treated ($n = 1580$) and non-PCI-treated ($n = 5052$) patients in the EPHESUS trial are shown in Table 1. The PCI-treated patients were younger, and had better renal function, fewer co-morbidities, and higher prescription of evidence-based medication including beta-blockers, ACE inhibitors, and statins. Cardiovascular mortality was significantly lower in PCI-treated patients as compared with non-PCI-treated patients (7% vs. 16%, $P < 0.0001$), but the incidence of non-fatal events was similar in the two groups (Table 1).

Eplerenone improves outcomes in both the PCI-treated and the non-PCI-treated cohort

Eplerenone treatment was similarly effective in reducing mortality, cardiovascular events, and hospitalization in both PCI-treated and non-PCI-treated AMI patients with LVSD (Figure 1). On multivariate Cox regression analysis, with eplerenone and PCI as co-variables, both PCI and eplerenone administration were independently associated with better clinical outcomes (Table 2).

Baseline characteristics of the EPHESUS PCI cohort

The baseline characteristics of the EPHESUS PCI cohort treated with placebo or eplerenone were similar (Table 3). The mean age of the patients was 60 years, with 77% male patients. The two groups were similar in terms of age, gender, ethnicity, LV function, co-morbidities, and prescription of medication at discharge (Table 3). There was no significant difference in adverse events between the eplerenone- or placebo-treated EPHESUS PCI cohort: hyperkalaemia (2.4% vs. 3.3%, $P = 0.33$), hypokalaemia (1.7% vs. 0.8%, $P = 0.10$), gynaecomastia in men (0.6% vs. 0.6%, $P = 1.00$), and renal dysfunction (3.2% vs. 2.0%, $P = 0.13$).

Eplerenone does not affect PCI-related clinical outcomes

Eplerenone, compared with placebo, in the PCI cohort did not impact upon the incidence of sudden cardiac death, non-fatal MI, or stroke (Table 4). Although there was a statistically significant reduction in events related to heart failure and hospital admissions, there was no effect upon PCI-related adverse outcomes, including recurrence of angina pectoris, unstable angina, AMI, or need for further revascularization (Table 4).

Discussion

This substudy analysis of the EPHESUS trial has highlighted that eplerenone is effective in patients with AMI and LVSD whether

Table 1 Baseline comparison of PCI-treated and non-PCI-treated EPHESUS patients

Characteristic	Non-PCI treated, n = 5052 (76%)		PCI treated, n = 1580 (24%)		P-value
	n	Mean ± SD or n (%)	n	Mean ± SD or n (%)	
Demographics					
Age (years)	5052	65 ± 11	1580	61 ± 12	<0.0001
Caucasian		4589 (91%)		1395 (88%)	0.003
Male		3491 (69%)		1223 (77%)	<0.0001
Time from AMI to randomization (days)	5051	7.4 ± 2.9	1580	6.7 ± 3.0	<0.0001
Clinical					
Blood pressure (mmHg)					
Systolic	5050	120 ± 17	1580	116 ± 16	<0.0001
Diastolic	5050	73 ± 11	1580	70 ± 11	<0.0001
LV ejection fraction (%)	5041	33 ± 6	1576	33 ± 6	0.85
Previous hospitalization for HF		452 (9%)		60 (4%)	<0.0001
Symptoms of heart failure		4347 (87%)		1232 (79%)	<0.0001
Potassium (mmol/L)	5029	4.28 ± 0.45	1566	4.21 ± 0.43	<0.0001
Creatinine (μmol/L)	5026	102 ± 29	1558	95 ± 25	<0.0001
eGFR (MDRD, mL/min/1.73 m ²)	4888	67 ± 19	1493	74 ± 19	<0.0001
Medical history	5052		1580		
Previous MI		1467 (29%)		336 (21%)	<0.0001
Diabetes mellitus		1658 (33%)		484 (31%)	0.10
Heart failure		853 (17%)		122 (8%)	<0.0001
Hypertension		3159 (63%)		848 (54%)	<0.0001
Medications	5052		1580		
ACEI/ARB		4318 (85%)		1433 (91%)	< 0.0001
Beta-blockers		3641 (72%)		1320 (84%)	< 0.0001
Diuretics		3172 (63%)		812 (51%)	< 0.0001
Aspirin		4377 (87%)		1493 (94%)	< 0.0001
Anticoagulants		934 (18%)		174 (11%)	< 0.0001
Statins		2026 (40%)		1069 (68%)	< 0.0001
Outcomes	5052		1580		
Cardiovascular mortality		784 (16%)		106 (7%)	<0.0001
Non-fatal events (ACS/revascularization)		1073 (21%)		339 (21%)	0.85

ACEI, ACE inhibitor; ACS, acute coronary syndrome; AMI, acute myocardial infarction; eGFR, estimated glomerular filtration rate [four-variable Modification of Diabetes in Renal Disease (MDRD) formula]; HF, heart failure; PCI, percutaneous coronary intervention.

treated with or without PCI. However, there is no evidence that eplerenone has any impact upon PCI-related outcomes including recurrence of angina, AMI, and repeat revascularization.

Eplerenone improved outcomes in both the PCI-treated and non-PCI-treated population

Mineralocorticoid receptor antagonists are strongly recommended for patients with LVSD and symptoms of heart failure,^{12,13} based on morbidity and mortality benefits seen in three landmark trials: RALES, EPHESUS, and EMPHASIS-HF.^{1,14,15} The RALES trial consisted of patients with advanced heart failure, the EPHESUS trial included patients with LVSD after AMI, and the EMPHASIS-HF trial enrolled patients with heart failure and mild (NYHA class

II) symptoms.^{1,14–16} In RALES, spironolactone produced 30% relative reduction in mortality during an average follow-up of 24 months.¹⁴ In EMPHASIS-HF, eplerenone was associated with a 24% reduction in cardiovascular death and a 42% reduction in hospitalization for heart failure.¹⁵ Only the EPHESUS trial specifically examined patients with LVSD after an AMI and showed that eplerenone produced 15% reduction in all-cause mortality, 17% reduction in cardiovascular mortality, and 21% reduction in sudden cardiac death.¹ In contemporary practice, most patients with AMI receive PCI treatment.⁴ Our data show that eplerenone confers similar benefit in patients with AMI and LVSD whether or not they are treated with PCI. The outcomes observed in the PCI subgroup were similar to those reported in the full EPHESUS cohort.¹ Our analysis shows that PCI itself was associated with a major reduction in adverse outcomes in AMI patients, consistent




Figure 1 Outcomes in PCI-treated and non-PCI-treated EPHEsus patients. Outcomes for both cohorts are shown as the hazard ratio (HR) with the 95% confidence interval (CI). The 95% CI for all HR values between PCI-treated and non-PCI-treated EPHEsus patients overlap with each other (shown in grey shade). CV, cardiovascular; PCI, percutaneous coronary intervention.

Table 2 Association of outcomes with eplerenone and PCI in EPHEsus patients in multivariable Cox regression analysis with eplerenone and PCI as covariables

	Eplerenone, yes vs. no		PCI, yes vs. no	
	HR (95% CI)	P-value	HR (95% CI)	P-value
Primary outcomes				
All-cause death	0.85 (0.75–0.96)	0.008	0.42 (0.35–0.51)	<0.0001
Cardiovascular death or hospitalization	0.87 (0.80–0.95)	0.003	0.62 (0.55–0.70)	<0.0001
Secondary outcomes				
All-cause death or any hospitalization	0.92 (0.86–0.99)	0.017	0.91 (0.84–0.99)	0.022
Cardiovascular death	0.83 (0.73–0.95)	0.005	0.41 (0.33–0.50)	<0.0001
Sudden death from cardiac causes	0.79 (0.64–0.97)	0.027	0.49 (0.36–0.65)	<0.0001
Acute myocardial infarction	0.82 (0.61–1.11)	0.20	0.34 (0.20–0.56)	<0.0001
Heart failure	0.80 (0.62–1.04)	0.099	0.35 (0.23–0.53)	<0.0001
Stroke	0.91 (0.54–1.56)	0.73	0.37 (0.16–0.87)	0.023
Any hospitalization	0.95 (0.89–1.02)	0.18	1.00 (0.92–1.09)	1.00
Cardiovascular hospitalization	0.91 (0.82–1.02)	0.098	0.73 (0.64–0.84)	<0.0001
Acute myocardial infarction	0.96 (0.80–1.16)	0.67	0.81 (0.65–1.02)	0.069
Heart failure	0.86 (0.75–0.98)	0.029	0.67 (0.56–0.80)	<0.0001
Stroke	1.18 (0.87–1.62)	0.29	0.40 (0.25–0.64)	0.0002
Ventricular arrhythmia	0.95 (0.65–1.39)	0.79	0.62 (0.37–1.03)	0.066
Acute myocardial infarction or unstable angina	1.01 (0.89–1.14)	0.92	0.96 (0.83–1.11)	0.60

CI, confidence interval; HR, hazard ratio; PCI, percutaneous coronary intervention.

Table 3 Baseline comparison of placebo- or eplerenone-treated patients in the EPHEUS-PCI group

Characteristic	Placebo, n = 781 (49%)		Eplerenone, n = 799 (51%)		P-value
	n	Mean $\pm$ SD or n (%)	n	Mean $\pm$ SD or n (%)	
Age (years)	781	61 $\pm$ 12	799	60 $\pm$ 11	0.05
Caucasian		688 (88%)		707 (88%)	0.81
Male gender		591 (76%)		632 (79%)	0.10
Blood pressure (mmHg)					
Systolic	781	116 $\pm$ 16	799	115 $\pm$ 15	0.43
Diastolic	781	70 $\pm$ 11	799	70 $\pm$ 10	0.65
LV ejection fraction (%)	779	33 $\pm$ 6	797	33 $\pm$ 6	0.83
Time from AMI to randomization (days)	781	6.6 $\pm$ 3.0	799	6.8 $\pm$ 3.0	0.23
Previous hospitalization for HF		34 (4%)		26 (3%)	0.25
Symptoms of heart failure		615 (79%)		617 (78%)	0.37
Potassium (mmol/L)	771	4.21 $\pm$ 0.43	795	4.21 $\pm$ 0.43	0.85
Creatinine (μ mol/L)	768	94 $\pm$ 25	790	95 $\pm$ 25	0.23
eGFR (MDRD, mL/min/1.73 m ²)	737	74 $\pm$ 19	756	73 $\pm$ 19	0.45
Medical history	781		799		
Previous MI		161 (21%)		175 (22%)	0.53
Diabetes mellitus		240 (31%)		244 (31%)	0.93
Heart failure		66 (8%)		56 (7%)	0.28
Hypertension		423 (54%)		425 (53%)	0.70
Medications	781		799		
ACEI/ARB		708 (91%)		725 (91%)	0.95
Beta-blockers		648 (83%)		672 (84%)	0.54
Diuretics		406 (52%)		406 (51%)	0.64
Aspirin		739 (95%)		754 (94%)	0.82
Anticoagulant agents		82 (10%)		92 (12%)	0.52
Statins		537 (69%)		532 (67%)	0.36

Symptoms of heart failure: Killip's class > I.

ACEI, ACE inhibitor; AMI, acute myocardial infarction; eGFR, estimated glomerular filtration rate [four-variable Modification of Diabetes in Renal Disease (MDRD) formula]; HF, heart failure; PCI, percutaneous coronary intervention.

Table 4 Effect of eplerenone on angioplasty-related outcomes in the EPHEUS-PCI cohort

	Eplerenone (n = 799)	Placebo (n = 781)	HR (95% CI)	P-value
Repeat revascularization (PCI/CABG)	51 (6%)	44 (6%)	1.13 (0.75–1.69)	0.56
Angina pectoris	24 (3%)	24 (3%)	0.97 (0.55–1.70)	0.90
Unstable angina	84 (11%)	82 (10%)	1.00 (0.73–1.35)	0.98
Acute myocardial infarction	45 (6%)	52 (7%)	0.83 (0.56–1.24)	0.37
Sudden cardiac death	21 (3%)	30 (4%)	0.67 (0.39–1.17)	0.16
Heart failure	60 (8%)	86 (11%)	0.66 (0.47–0.91)	0.012
Cardiovascular hospitalization	108 (14%)	140 (18%)	0.72 (0.56–0.93)	0.011

CABG, coronary artery bypass grafting; CI, confidence interval; HR, hazard ratio.

with numerous other reports.^{17,18} However, eplerenone administration was also an independent factor associated with improved outcomes, emphasizing the importance of MR antagonism in LVSD after AMI, including patients treated with primary PCI. Therefore, prescribing eplerenone to patients with AMI and LVSD remains beneficial in patients treated with and without PCI.

Whilst the EPHEUS trial included patients with all forms of AMI [unstable angina, non-ST segment elevation myocardial

infarction (NSTEMI), and STEMI] and there was a delay in commencing eplerenone, a recently reported clinical trial (REMINDER: impact of eplerenone on cardiovascular outcomes in patients post myocardial infarction, NCT-01176968) has shown that early use of eplerenone can improve the outcome of patients presenting with acute STEMI without heart failure.¹⁹ Over 1000 patients were randomized (n = 506/group) to initiation of eplerenone or placebo within 12–24 h of STEMI. Eplerenone 25 or 50 mg/day (89% patients received 50 mg/day) produced a significant reduction

(HR 0.58, 95% CI 0.45–0.75, $P < 0.0001$) in the primary endpoint (a composite of cardiovascular mortality, ventricular arrhythmia, clinical or subclinical heart failure at 1 month) at a mean follow-up of 10.5 months.¹⁹ ALBATROSS (Aldosterone blockade early after acute myocardial infarction; NCT-01059136) is an ongoing multi-centre, open-labelled, randomized trial to assess the effects of MR blockade with a 200 mg i.v. bolus of potassium canrenoate followed by 25 mg/day spironolactone for 6 months in 1600 patients with STEMI or high risk NSTEMI.²⁰

Eplerenone did not affect PCI-related outcomes

Mineralocorticoid receptors are widely distributed in the vascular system, including endothelial cells, endothelial progenitor cells (EPCs), vascular smooth muscle cells (VSMCs), and intact vessels.^{8,21,22} These receptors also influence vascular function and remodelling.^{6,23} MR activation has been shown to increase proliferation of VSMCs⁶ and to impair differentiation and proliferation of EPCs.⁷ These characteristics suggest that MR activation is a potential stimulus for neointimal proliferation and restenosis. In humans, elevated baseline plasma aldosterone levels directly correlate with risk of restenosis²⁴ and inversely correlate with circulating EPCs and endothelial colony-forming units.⁸ Mineralocorticoid receptor antagonists reduce atherosclerosis,^{23,25} prevent adverse vascular remodelling,²⁶ and attenuate post-angioplasty neointimal proliferation in experimental animals.^{9,11,27}

Spironolactone has been shown to inhibit neointimal proliferation after balloon angioplasty of iliac arteries in rabbits.²⁷ However, these results could not be reproduced in a porcine coronary angioplasty model,^{9,28} which is the gold standard pre-clinical model for restenosis. Additionally, a single-centre clinical trial failed to show any effect of spironolactone in reducing restenosis.²⁹ Nevertheless, promising experimental data have been obtained with eplerenone.^{9–11} Our data suggest that, like spironolactone, eplerenone does not appear to affect PCI-related clinical outcomes. Therefore, eplerenone cannot currently be recommended for patients undergoing PCI without co-existing AMI or LVSD.

Study limitations

This study has several limitations. It is a post-hoc analysis with inherent shortcomings of any such study. Patients in the EPHEUS PCI cohort did not have routine angiographic follow-up to document any effect of eplerenone on angiographic restenosis; however, our assessment of clinical events is perhaps more relevant than angiographic outcomes.

Conclusion

The beneficial effects of eplerenone on heart failure events and hospitalization seen in the EPHEUS trial are similar for both PCI-treated and non-PCI-treated AMI patients with LVSD. There is no evidence that eplerenone reduces the risk of recurrent ischaemia-related events including recurrence of angina or the

need for repeat revascularization. Use of eplerenone is strongly recommended in AMI patients with LVSD.

Funding

The EPHEUS trial was funded by Pharmacia/Pfizer.

Conflict of interest: I.S. has received grants and honoraria from Novartis, Pfizer, Vifor, and Jansen. B.P. has received honoraria from Pfizer and serves on the advisory boards for Pfizer and Novartis. F.Z. has received honoraria from Pfizer, Novartis, Roche, Servier, AstraZeneca, and Takeda. All other authors have no conflicts to declare.

References

- Pitt B, Remme W, Zannad F, Neaton J, Martinez F, Roniker B, Bittman R, Hurley S, Kleiman J, Gatlin M. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 2003;**348**:1309–1321.
- Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, Jessup M, Konstam MA, Mancini DM, Michl K, Oates JA, Rahko PS, Silver MA, Stevenson LW, Yancy CW. 2009 Focused update incorporated into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults. A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines developed in collaboration with the International Society for Heart and Lung Transplantation. *J Am Coll Cardiol* 2009;**53**:e1–e90.
- Dickstein K, Cohen-Solal A, Filippatos G, McMurray JJ, Ponikowski P, Poole-Wilson PA, Stromberg A, van Veldhuisen DJ, Atar D, Hoes AV, Keren A, Mebazaa A, Nieminen M, Priori SG, Swedberg K. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the diagnosis and treatment of acute and chronic heart failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur J Heart Fail* 2008;**10**:933–989.
- O'Gara PT, Kushner FG, Ascheim DD, Casey DE, Chung MK, de Lemos JA, Ettinger SM, Fang JC, Fesmire FM, Franklin BA, Granger CB, Krumholz HM, Linderbaum JA, Morrow DA, Newby LK, Ornato JP, Ou N, Radford MJ, Tamis-Holland JE, Tommaso CL, Tracy CM, Woo YJ, Zhao DX. 2013 ACCF/AHA Guideline for the Management of ST-Elevation Myocardial Infarction. A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;**61**:e78–e140.
- Wijns W, Kolh P, Danchin N, Di Mario C, Falk V, Folliquet T, Garg S, Huber K, James S, Knuuti J, Lopez-Sendon J, Marco J, Menicanti L, Ostojic M, Piepoli MF, Pirlot C, Pomar JL, Reifart N, Ribichini FL, Schalij MJ, Sergeant P, Serruys PV, Silber S, Sousa Uva M, Taggart D. Guidelines on myocardial revascularization. *Eur Heart J* 2010;**31**:2501–2555.
- Ishizawa K, Izawa Y, Ito H, Miaki C, Miyata K, Fujita Y, Kanematsu Y, Tsuchiya K, Tamaki T, Nishiyama A, Yoshizumi M. Aldosterone stimulates vascular smooth muscle cell proliferation via big mitogen-activated protein kinase 1 activation. *Hypertension* 2005;**46**:1046–1052.
- Thum T, Schmitter K, Fleissner F, Wiebking V, Dietrich B, Widder JD, Jazbutyte V, Hahner S, Ertl G, Bauersachs J. Impairment of endothelial progenitor cell function and vascularization capacity by aldosterone in mice and humans. *Eur Heart J* 2011;**32**:1275–1286.
- Wu VC, Lo SC, Chen YL, Huang PH, Tsai CT, Liang CJ, Kuo CC, Kuo YS, Lee BC, Wu EL, Lin YH, Sun YY, Lin SL, Chen JW, Lin SJ, Wu KD. Endothelial progenitor cells in primary aldosteronism: a biomarker of severity for aldosterone vasculopathy and prognosis. *J Clin Endocrinol Metab* 2011;**96**:3175–3183.
- Ward MR, Kanellakis P, Ramsey D, Funder J, Bobik A. Eplerenone suppresses constrictive remodeling and collagen accumulation after angioplasty in porcine coronary arteries. *Circulation* 2001;**104**:467–472.
- Wakabayashi K, Suzuki H, Sato T, Iso Y, Katagiri T, Takeyama Y. Eplerenone suppresses neointimal formation after coronary stent implantation in swine. *Int J Cardiol* 2006;**107**:260–266.
- Iqbal J, Macdonald LJ, Low L, Seckl JR, Yau CW, Walker BR, Hadoke PV. Contribution of endogenous glucocorticoids and their intravascular metabolism by 11 β -HSDs to postangioplasty neointimal proliferation in mice. *Endocrinology* 2012;**153**:5896–5905.

12. Jessup M, Abraham WT, Casey DE, Feldman AM, Francis GS, Ganiats TG, Konstam MA, Mancini DM, Rahko PS, Silver MA, Stevenson LW, Yancy CW. 2009 focused update: ACCF/AHA Guidelines for the Diagnosis and Management of Heart Failure in Adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the International Society for Heart and Lung Transplantation. *Circulation* 2009;**119**:1977–2016.
13. McMurray JJ, Adamopoulos S, Anker SD, Auricchio A, Bohm M, Dickstein K, Falk V, Filippatos G, Fonseca C, Gomez-Sanchez MA, Jaarsma T, Kober L, Lip GY, Maggioni AP, Parkhomenko A, Pieske BM, Popescu BA, Ronnevik PK, Rutten FH, Schwitler J, Seferovic P, Stepinska J, Trindade PT, Voors AA, Zannad F, Zeiher A, Bax JJ, Baumgartner H, Ceconi C, Dean V, Deaton C, Fagard R, Funck-Brentano C, Hasdai D, Hoes A, Kirchhof P, Knuuti J, Kolh P, McDonagh T, Moulin C, Reiner Z, Sechtem U, Sirnes PA, Tendera M, Torbicki A, Vahanian A, Windecker S, Bonnet LA, Avraamides P, Ben Lamin HA, Brignole M, Coca A, Cowburn P, Dargie H, Elliott P, Flachskampf FA, Guida GF, Hardman S, Jung B, Merkely B, Mueller C, Nanas JN, Nielsen OV, Orn S, Parissis JT, Ponikowski P. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2012;**14**:803–869.
14. Pitt B, Zannad F, Remme WJ, Cody R, Castaigne A, Perez A, Palensky J, Wittes J. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N Engl J Med* 1999;**341**:709–717.
15. Zannad F, McMurray JJ, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pocock SJ, Pitt B. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med* 2011;**364**:11–21.
16. Iqbal J, Parviz Y, Pitt B, Newell-Price J, Al-Mohammad A, Zannad F. Selection of a mineralocorticoid receptor antagonist for patients with hypertension or heart failure. *Eur J Heart Fail* 2014;**16**:143–150.
17. Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet* 2003;**361**:13–20.
18. Fox KA, Poole-Wilson PA, Henderson RA, Clayton TC, Chamberlain DA, Shaw TR, Wheatley DJ, Pocock SJ. Interventional versus conservative treatment for patients with unstable angina or non-ST-elevation myocardial infarction: the British Heart Foundation RITA 3 randomised trial. *Randomized Intervention Trial of unstable Angina. Lancet* 2002;**360**:743–751.
19. Montalescot G, Pitt B, Lopez-de-Sa E, Hamm CW, Flather M, Verheugt F, Shi H, Turgonyi E, Orri M, Vincent J, Zannad F. Early eplerenone treatment in patients with acute ST-elevation myocardial infarction without heart failure. ACC, San Francisco, 10 March, 2013.
20. Beygui F, Vicaut E, Ecollan P, Machecourt J, Van Belle E, Zannad F, Montalescot G. Rationale for an early aldosterone blockade in acute myocardial infarction and design of the ALBATROSS trial. *Am Heart J* 2010;**160**:642–648.
21. Hadoke PW, Iqbal J, Walker BR. Therapeutic manipulation of glucocorticoid metabolism in cardiovascular disease. *Br J Pharmacol* 2009;**156**:689–712.
22. Kayes-Wandover KM, White PC. Steroidogenic enzyme gene expression in the human heart. *J Clin Endocrinol Metab* 2000;**85**:2519–2525.
23. Suzuki J, Iwai M, Mogi M, Oshita A, Yoshii T, Higaki J, Horiuchi M. Eplerenone with valsartan effectively reduces atherosclerotic lesion by attenuation of oxidative stress and inflammation. *Arterioscler Thromb Vasc Biol* 2006;**26**:917–921.
24. Amano T, Matsubara T, Izawa H, Torigoe M, Yoshida T, Hamaguchi Y, Ishii H, Miura M, Hayashi Y, Ogawa Y, Murohara T. Impact of plasma aldosterone levels for prediction of in-stent restenosis. *Am J Cardiol* 2006;**97**:785–788.
25. Keidar S, Hayek T, Kaplan M, Pavlotzky E, Hamoud S, Coleman R, Aviram M. Effect of eplerenone, a selective aldosterone blocker, on blood pressure, serum and macrophage oxidative stress, and atherosclerosis in apolipoprotein E-deficient mice. *J Cardiovasc Pharmacol* 2003;**41**:955–963.
26. Endemann DH, Touyz RM, Iglarz M, Savoia C, Schiffrin EL. Eplerenone prevents salt-induced vascular remodeling and cardiac fibrosis in stroke-prone spontaneously hypertensive rats. *Hypertension* 2004;**43**:1252–1257.
27. Van Belle E, Bauters C, Wernert N, Hamon M, McFadden EP, Racadot A, Dupuis B, Lablanche JM, Bertrand ME. Neointimal thickening after balloon denudation is enhanced by aldosterone and inhibited by spironolactone, and aldosterone antagonist. *Cardiovasc Res* 1995;**29**:27–32.
28. Bacchetta MD, Salemi A, Milla F, Hong MK, Tio F, Zhou Y, Chen R, Southard E, Lee LY, Mack CA, Krieger KH, Isom OW, Ko W, Borer JS, Catanzaro DF. Low-dose spironolactone: effects on artery-to-artery vein grafts and percutaneous coronary intervention sites. *Am J Ther* 2009;**16**:204–214.
29. Kursaklioglu H, Iyisoy A, Amasyali B, Celik T, Ozturk C, Kose S, Isik E. Spironolactone does not prevent restenosis after coronary stenting in humans. *Ann Acad Med Singapore* 2004;**33**:769–774.